



Rearrangement of Molecular Complexes

Rainer Streubel and Gerd von Frantzius

published in

NIC Symposium 2006,
G. Münster, D. Wolf, M. Kremer (Editors),
John von Neumann Institute for Computing, Jülich,
NIC Series, Vol. 32, ISBN 3-00-017351-X, pp. 83-90, 2006.

© 2006 by John von Neumann Institute for Computing

Permission to make digital or hard copies of portions of this work for personal or classroom use is granted provided that the copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise requires prior specific permission by the publisher mentioned above.

<http://www.fz-juelich.de/nic-series/volume32>


Rearrangement of Molecular Complexes

Rainer Streubel and Gerd von Frantzius

Institut für Anorganische Chemie
Gerhard-Domagk-Str. 1, Universität Bonn, Germany
E-mail: {r.streubel, gfrantzi}@uni-bonn.de

1 Introduction

Experimental and computational investigations of rearrangement and decomposition pathways of molecular phosphine transition-metal complexes are of fundamental interest in coordination and organometallic chemistry and in catalysis. One of the central topics of our research group is the chemistry of *2H*-azaphosphirene complexes **1** and the generation, reaction and/or rearrangement of reactive intermediates **2** (i) (so called phosphinidene complexes) and **3** (nitrilium phosphanylide complexes) (ii) from precursor **1**¹⁻³. Recently, we had discovered that electrophilic terminal phosphinidene tungsten complexes **2** with bulky substituents at phosphorus (R = C₅Me₅ or CH(SiMe₃)₂) do not dimerize in the absence of trapping reagents⁴. Therefore we became interested in their fate. The computational studies comprise a density functional method-based study of the hypersurfaces of phosphinidene complexes **2** and nitrilium phosphanylide complexes **1** as well as a characterization of bonding situations by the method of the compliance matrix; probing and establishing this rarely used method - especially in the field of transition metal complexes - makes it a goal in itself and will be discussed below.



Scheme 1. Rearrangement and decomposition of *2H*-azaphosphirene complexes **1**.

2 Computational Methods

All optimizations and frequency calculations were performed using GAUSSIAN 03 RevB.03⁵ on the IBM Regatta p690 cluster (JUMP) of the John von Neumann Institute for Computing (NIC) at the Forschungszentrum Jülich. The standard method throughout this work is B3LYP/6-311G(d,p) combined with an effective core potential description of tungsten using the Los Alamos LanL2DZ (for short: B3LYP/6-311**/LanL2DZ)⁶. Stationary points have been characterized by analytical second derivatives (the Hessian) with respect to redundant cartesian coordinates. Transition states (one imaginary frequency)

were identified by a reaction path following (IRC). The Hessian provided by the GAUSSIAN 03 calculation was transformed to non-redundant internal coordinates using Pulays INTC/FCTINT set of algorithms⁷. The inversion of the transformed Hessian to the compliance matrix was accomplished by standard methods.⁸ A typical production job, e.g. C₁₉H₂₄NO₅PSi₂W **1** (Scheme 1) with 242 electrons, uses more than 1000 Gaussian-basis functions and thus needs parallelized computations on fast processors, *which can only be provided by a computing centre like the John von Neumann Institute for Computing (NIC)*.

3 Usage and Interpretation of the Compliance Matrix

The physical model behind force constants and thus compliance constants is a spring model: if the molecule at equilibrium geometry is distorted by a vibrational movement the various internal coordinates (modeled by springs) interact according to the molecular force field. In the harmonic approximation to vibrational theory the molecular hypersurface describing a vibrational movement is locally approximated by a quadratic form (H_{ik}) in the displacements of the internal coordinates (bonds, angles, dihedrals and linear combinations thereof) from their equilibrium values

$$2V = \Delta x^t (H_{ik}) \Delta x, \quad (1)$$

where (H_{ik}) is the matrix of second derivatives (Hessian) at a stationary point, the diagonal elements H_{kk} are the force constants⁹. Equivalently the change in potential energy during a vibration is described by a quadratic form in the forces (force displacements Δf to be precise) instilled in the coordinates upon distortion from equilibrium geometry

$$2V = \Delta f^t (C_{ik}) \Delta f, \quad (2)$$


where (C_{ik}) = (H_{ik})⁻¹ is the compliance matrix (inverse Hessian) at a stationary point, the diagonal elements C_{kk} of which are the compliance constants (in [$\text{\AA}/\text{mdyn}$] for bond stretchings and in [rad/mdyn] for angle bendings); the off-diagonal elements are associated with the couplings of the coordinates. While in the spring model force constants H_{kk} describe the stiffness (resistance against distortion) the compliance constants C_{kk} are associated with the compliance of coordinate k ¹⁰.

4 Experimental and Theoretical Results

4.1 Formation of a Nitrilium Phosphanylide Complex

From experimental investigations of the thermolysis of 2*H*-azaphosphirene complex **1a**¹¹ the existence of nitrilium phosphanylide complex **3a** (Scheme 1, path ii), R = CH(SiMe₃)₂, Ar = Ph) as a reactive intermediate has been concluded, although, due to its assumed short lifetime, there was no spectroscopic evidence for the intermediacy of **3a**, so far. The DFT-calculations on the hypersurface of **1a** unambiguously showed that nitrilium phosphanylide complex **3a** is an isomer; the formation of which via a still unknown transition state proceeds endergonically by 19 kJ/mol. A comparison of selected experimental (X-ray crystal structure) and calculated bond lengths of **1a** as well as calculated bond lengths of **3a** can be taken from table 1. There is satisfying agreement between experiment and theory concerning the geometry of **1a**. It is noteworthy, that in **3a** the almost linear benzonitrile unit

(angle NCC in **1a**: 135 degree) is attached to a strongly pyramidal phosphorus (sum of bond angles at P: 326 degree); the compliance constant of the P-N contact (0.552 Å/mdyn) provides a likely description of **3a** as a N-P donor adduct of benzonitrile to phosphinidene complex **2b** (see Scheme 2, Figure 2 below).




	1a Exp. [Å], [°]	1a Calc. [Å], [°]	3a Calc. [Å], [°]
W-P	2.470(2)	2.551	2.628
P-C(N)	1.759(5)	1.781	-
P-C(H)	1.808(5)	1.849	1.892
P-N	1.795(4)	1.821	1.752
C-N	1.272(7)	1.271	1.165
$\Sigma \angle (\text{P})$			326

Figure 1. (hydrogens except PCH are omitted) Table 1

4.2 Formation of a *P*-SiMe₃-Substituted η^1 -*E*-Phosphaalkene Complex¹²

As experimentally shown heating diluted ortho-xylene solutions of *2H*-azaphosphirene complex **1a** afforded almost quantitatively and stereoselectively the *P*-SiMe₃-substituted η^1 -*E*-phosphaalkene complex **4a** (Scheme 2). The rearrangement of the thermally generated short-lived phosphinidene complex [(OC)₅W{PCH(SiMe₃)₂}] **2** to complex **4** was completely unexpected; we assume that a 1,2-(C-P)-trimethylsilyl shift takes place in this case. Although the chemistry of short-lived electrophilic terminal phosphinidene tungsten complexes¹³ has received increased attention during the last years, partially because of the versatility of *2H*-azaphosphirene complexes,¹⁴ only a single example of a rearrangement yielding a *P*-Cl-substituted η^1 -phosphaalkene complex - via a 1,2-chlorine shift - has been reported, so far¹⁵.



Scheme 2. Rearrangement of transient phosphinidene complexes **2a,b** into **4a,b**.

DFT calculations in order to study the rearrangement of pentacarbonyl-tungsten phosphinidene complexes **2a** (*anti*-periplanar) and **2b** (*syn*-periplanar) to the corresponding phosphaalkene complexes **4a** and **4b** revealed that the formation of the C-P double bonds proceeds via transition state complexes **TSa** and **TSb** (Figure 2; hydrogens except P=CH are omitted).




Figure 2. Phosphinidene complexes **2a,b**, transition states **TSa,b** and phosphalkene complexes **4a,b**.

In consequence, this rearrangement represents an interesting and unique example of an intramolecular 1,2-silyl shift in phosphinidene complex chemistry. Relative energies of the 1,2-silyl migration of **2a,b** to corresponding **4a,b** can be taken from figure 3. *Anti*-periplanar phosphinidene complex **2a** rearranges via transition state **TSa** to the *E*-





Figure 3. Energy profile of the 1,2-(C to P)-silyl migrations in **2a** and **2b**.

phosphalkene complex **4a**. C_1 symmetric phosphinidenes **2a,b** exhibit a long and a short C-Si bond; the former is the one involved in forming the three-membered ring of the transition state which contains a five-coordinated silicon center. From an estimated singlet-triplet gap of about 40 kJ/mol under thermal reaction conditions phosphinidene complex **2a** can choose from two concurrent pathways: either undergo a fast reaction from the triplet excited state or - in the absence of suitable reaction partners - rearrange to *E*-phosphalkene complex **4a**; the reverse 1,2-silyl shift is hindered by a substantial barrier of about 144 kJ/mol. These findings agree well with experimental results where reaction conditions

could be optimized to synthesize pentacarbonyltungsten phosphalkene complex **4a** from the 2*H*-azaphosphirene complex **1a**. Apart from a smaller barrier (54 kJ/mol **2a** to **4a** vs. 74 kJ/mol for **2b** to **4b**) the major difference between **TSa** and **TSb** can be seen by comparison of the P-Si and C-Si bond lengths involved in the transition state. While **TSa** is early (according to the C-Si bond lengths) in **TSb** the position of the silicon center is almost in between the originating carbon and the phosphorus.


4.3 Further Local Minima on the Hypersurface of [(OC)₅W{PCH(SiMe₃)₂}]

So far at least six further local minima and a transition state have been located; four of them are shown in scheme 3 and figure 4.



Scheme 3.

Due to the CH(SiMe₃)₂ substituent at phosphorus phosphinidene complexes **2a,b** can also undergo a 1,2-*H*-shift to yield phosphalkene complexes **5a,b** and, by a still unknown transition state, highly unusual complex **6a**. The nature of the related transition-state complex **6b** (figure 4) remains to be clarified since reaction path following calculations were not conclusive yet. There is no automated way to find isomers of a given sum formula. From the literature¹⁶ the ability of phosphinidene complexes to undergo intramolecular C-H insertions is known. Thus we could locate corresponding isomers **7a,b** (Scheme 4, Figure 4; only selected hydrogens are shown for clarity), which are thermodynamically more stable than phosphinidene complexes **2a,b** by 100 kJ/mol and even more stable than the phosphalkene complexes **4** by 30 kJ/mol (**7b**) and by 20 kJ/mol (**7a**) respectively.



Scheme 4. Intramolecular C-H insertion reaction of phosphinidene complexes **2a,b** to yield **7a,b**.

5 Use of Compliance Constants for the Description of Bonding Situations

A compliance constant is the displacement of a bond due to the application of a unit force on that bond including reorganization. That means, a higher numerical value is connected




Figure 4.

with a weaker bond. Introduced to vibrational theory by Decius¹⁷ and others¹⁸ experimental compliance fields obtained by Jones and Swanson from vibrational data clarified the bonding forces in metal cyanides and carbonyls¹⁹, while Williams used compliance matrices for the description of chemical reactivity²⁰. Calculations of full compliance fields have been used to investigate Ga-Ga²¹ and Si-C²² multiple bonds, the metal-metal bond character in homoleptic transition metal carbonyls²³, hydrogen bonds in Watson-Crick base pairs²⁴ and polyphosphorus compounds²⁵. Recently, Andreoni and coworkers used compliance constants plus Car-Parinello molecular dynamic simulations in order to analyze a proposed Si-Si triple bond²⁶. We could show recently the usefulness of compliance constants (diagonal elements of the compliance matrix) to assess the strength of a particular bond type (e.g. a tungsten-phosphorus bond) a set of reference compounds has to be calculated (Figure 5). From the statistic three different strengths of bonding interactions can be inferred: tungsten-phosphorus triple bonds range from 0.2-0.3 Å/mdyn while phosphinidene complexes of the type $[(OC)_5WR]$ seem to have a $W(CO)_5$ unit doubly bonded to phosphorus; ordinary W-P single bonds have compliance constants greater than 0.7 Å/mdyn.




Figure 5. Compliance constants of P-C bonded compounds²⁸.

By a similar procedure phosphorus-carbon triple (0.1 Å/mdyn), double (0.15-0.2 Å/mdyn) and single bonds (0.3-0.4 Å/mdyn) can be identified (Figure 6). For example the P-C compliance constant of phosphalkene complex $[(OC)_5W\{HP=C(SiMe_3)_2\}]$ (Scheme 3) falls

well within the range of a double bond (0.192 Å/mdyn).




Figure 6. Compliance constants of P-C bonded compounds²⁹.

Acknowledgments

We are grateful to the *Deutsche Forschungsgemeinschaft* and the *Fonds der Chemischen Industrie* for financial support; we thank the John von Neumann Institute for Computing (NIC) at the Forschungszentrum Jülich for providing computational resources.

References

1. Review on the chemistry of λ^3 -2H-azaphosphirene metal complexes: R. Streubel, *Coord. Chem. Rev.* **2002**, 227, 175.
2. Review on phospho-organic chemistry: F. Mathey, *Angew. Chem. Int. Ed.* **2003**, 42, 1578.
3. Reviews on phosphinidene complexes: K. Lammertsma, *Top. curr. Chem.* **2002**, 229, 35; F. Mathey, *Angew. Chem. Int. Ed.* **2003**, 42, 1586.
4. a) H. Wilkens, A. Ostrowski, J. Jeske, F. Ruthe, P. G. Jones, R. Streubel, *Organometallics* **1999**, 18, 5627; b) R. Streubel, U. Schiemann, N. Hoffmann, Y. Schiemann, P. G. Jones, D. Gudat, *Organometallics* **2000**, 19, 475.
5. Gaussian 03, Revision B.02/B.03, Gaussian, Inc., Wallingford CT, **2004**.
6. a) A. D. Becke, *J. Chem. Phys.* **1993**, 98, 5648. b) R. Krishnan, J. S. Binkley, R. Seeger, J. A. Pople, *J. Chem. Phys.* **1980**, 72, 650. c) A. D. McLean, G. S. Chandler, *J. Chem. Phys.* **1988**, 53, 1995. d) P. J. Hay, W. R. Wadt, *J. Chem. Phys.* **1988**, 82, 270.
7. G. Fogarasi, X. Zhou, P. W. Taylor, P. Pulay, *J. Am. Chem. Soc.* **1992**, 114, 8191.
8. W. H. Press, S. A. Teukolsky, W. T. Vetterling, B. P. Flannery, *Numerical Recipes in Fortran: The Art of Scientific Computing*, Cambridge University Press, Cambridge, **1994**.

9. E. B. Wilson, *J. Chem. Phys.* **1939**, 7, 1047.
10. L. H. Jones, B. I. Swanson, *Acc. Chem. Res.* **1976**, 9, 128.
11. R. Streubel, A. Ostrowski, S. Priemer, U. Rohde, J. Jeske, P. G. Jones, *Eur. J. Inorg. Chem.* **1998**, 257.
12. E. Ionescu, Gerd v. Frantzius, R. Streubel, *Manuscript in Preparation*.
13. Recent reviews: F. Mathey, N. G. Tran Huy, M. Marinetti, *Helv. Chim. Acta* **2001**, 84, 2938; K. Lammertsma, M. J. M. Vlaar, *Eur. J. Org. Chem.* **2002**, 1127.
14. a) R. Streubel, *Coord. Chem. Rev.* **2002**, 227, 172; ; b) R. Streubel, *Top. Curr. Chem.* **2002**, 223, 91.
15. B. Deschamps, F. Mathey, *J. Organomet. Chem.* **1988**, 354, 83 (Cu(I) catalyzed shift).
16. A. H. Cowley, A. R. Barron, *Acc. Chem. Res.* **1988**, 21, 81.
17. J. C. Decius, *J. Chem. Phys.* **1963**, 38, 241.
18. See: D. Papousek, J. Pliva, *Spectrochimica Acta* **1965**, 21, 1147.
19. L. H. Jones, B. I. Swanson, *Acc. Chem. Res.* **1976**, 9, 128.
20. I. H. Williams, *Chem. Phys. Lett.* **1982**, 88, 462.
21. J. Grunenberg, *J. Chem. Phys.* **2001**, 115, 6360.
22. a) J. Grunenberg, *Angew. Chem.* **2001**, 113, 4150. b) J. Grunenberg, R. Streubel, G. von Frantzius, W. Marten, *J. Chem. Phys.* **2003**, 119, 165.
23. Y. Cie, H. F. Schaefer III, *Z. Phys. Chem.* **2003**, 217, 189.
24. J. Grunenberg, *J. Am. Chem. Soc.* **2004**, 126, 1631.
25. S. Katsyuba, R. Schmutzler, J. Grunenberg, *J. Chem. Soc., Dalton Trans.* **2005**, 9, 1700.
26. C. A. Pignedoli, A. Curioni, W. Andreoni, *Chem. Phys. Chem.* **2005**, 6, 1795.
27. G. von Frantzius, R. Streubel, K. Brandhorst, J. Grunenberg, *Organometallics*, Web Release Date: 08-Dec- **2005**.
28. The complexes used for this calculation were first published by the following authors: **8**: R. Schrock, N.C. Zanetti, *Angew. Chem. Int. Ed. Engl.* **1995**, 34, 2044. **9**: [(OC)₅WP]⁺ (this work). **10**: [(OC)₂CpW=P^tBu₂] J. Klaus, W. Malisch, W. Reich, A. Meyer, U. Schubert, *Angew. Chem.* **1986**, 98, 103. **11-13**: K. Lammertsma, *Top. Curr. Chem.* **2003**, 229, 95. **2a**: [(OC)₅W{PCH(SiMe₃)₂}] (this work). **15**: G. Frenking, K. Wichmann, N. Frhlich, J. Grobe, W. Golla, D. Le Van, B. Krebs, M. Läge, *Organometallics* **2002**, 21, 2921. **17**: [(OC)₃CpW-P=C(SiMe₃)₂] D. Gudat, E. Niecke, W. Malisch, U. Hofmockel, S. Quashie, A. H. Cowley, A.M. Arif, B. Krebs, M. Dartmann, *J. Chem. Soc., Chem. Commun.* **1985**, 1687.
29. The complexes used for this calculation were first published by the following authors: **16**: This work. **14**: W. W. Schoeller, A. B. Rozhenko, S. Grigoleit, *Eur. J. Inorg. Chem.* **2001**, 2891. **4a**: [(OC)₅W{Me₃SiP=C(H)SiMe₃}] (this work). **17**: [(OC)₃CpW-P=C(SiMe₃)₂] D. Gudat, E. Niecke, W. Malisch, U. Hofmockel, S. Quashie, A. H. Cowley, A.M. Arif, B. Krebs, M. Dartmann, *J. Chem. Soc., Chem. Commun.* **1985**, 1687. **18**: M. I. Povolotskii, V. V. Negrebetskii, *J. Gen. Chem. USSR* **1988**, 58, 206. **12,13**: K. Lammertsma, *Top. Curr. Chem.* **2003**, 229, 95. **19**: H. Vahrenkamp, et al, *Z. Naturf. B.* **1985**, 40, 1250; F. Nief, F. Mercier, F. Mathey, *J. Organomet. Chem.* **1987**, 328, 349. **20**: K. Miqueu, J.-M. Sotiropoulos, G. Pfister-Guillouzo, V. Romanenko, *Eur. J. Inorg. Chem.* **2000**, 3, 477. **2b**: [(OC)₅W{PCH(SiMe₃)₂}] (this work).